


Kitsap County

Adults


Current Cigarette Smoking (1987-2002)



Current Smokeless Tobacco Use (1999-2001)

	<i>Entire adult population</i>	<i>Males only</i>
Kitsap County	1.4% (95%CI: 0.6,3.5)	3.0% (95%CI: 1.2,7.4)
Washington State	2.8% (95%CI: 2.5,3.2)	5.6% (95%CI: 4.9,6.4)


Youth



* = RATE NOT CALCULATED BECAUSE COUNTY DID NOT MEET MINIMUM PARTICIPATION REQUIREMENTS FOR THAT GRADE.

DATA SOURCES: (ADDITIONAL DOCUMENTATION INCLUDED IN APPENDIX A, WWW.DOH.WA.GOV/TOBACCO/DATA/COUNTY_PROFILES/PROFILES_2002/APNDXA.PDF)
 ADULT TOBACCO USE – WA. DEPT OF HEALTH, BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (BRFSS), 2002 PRELIMINARY ESTIMATE BASED ON FIRST SIX MONTHS OF YEAR; YOUTH TOBACCO USE – WA STATE SURVEY OF ADOLESCENT HEALTH BEHAVIORS (WSSAHB); POPULATION ESTIMATES – OFFICE OF FINANCIAL MGMT(OFM), YEAR 2000 CENSUS; MATERNAL SMOKING DURING PREGNANCY – WA BIRTH CERT SYSTEM; LUNG AND BRONCHUS CANCER – WA STATE CANCER REGISTRY; ESTIMATED TOTAL DEATHS – WA DEATH CERT SYSTEM.

Pregnant Women



Summary of the problem


- Exhibiting a sinuous pattern around Washington State as a whole, adult cigarette smoking is not significantly different from the state overall. Our most recent survey data showed that about 20.7% of adults in Kitsap County smoke cigarettes.
- Although adult smokeless tobacco use is lower than Washington State as a whole, it is not significantly different from the state overall.
- Youth rates could not be calculated because Kitsap County did not meet minimum participation requirements. More data collection is needed to describe Kitsap County youth specifically.
- Maternal smoking during pregnancy is significantly higher than the state average in each of the last eight years. In recent years, the trend is the same as the state rate. In 2001, about 16.4% of pregnant women in Kitsap County smoked during pregnancy, compared to 12.6% statewide.

People Affected in Kitsap County

- *Total Population: 231,969 - 18 and over: 169,905 (73%); under 18: 62,064 (27%)*
- *Average number of people diagnosed with Lung and Bronchus cancer each year: 51*
- *Infants born annually to mothers who smoke during pregnancy: 480*
- *Estimated total deaths due to tobacco use each year: 330*

Melinda Harmon
 Kitsap Health District
 109 Austin Drive
 Bremerton, WA 98312
 (360) 337-5224
 fax: (360) 337-5298
 email: harmon@health.co.kitsap.wa.us

website: <http://www.wa.gov/kitsaphealth>



TECHNICAL NOTES: TOTAL DEATHS DUE TO TOBACCO USE ARE ESTIMATED AS 19% OF ALL DEATHS (McGINNIS AND FOEGE 1993); "DUMBELL" BARS AROUND DATA POINTS FOR SURVEY DATA REPRESENT 95% CONFIDENCE INTERVALS – THE TRUE VALUE MAY BE ANYWHERE WITHIN THE BARS, BUT MAY NOT BE EXACTLY AT THE DATA POINT.

ADDITIONAL DOCUMENTATION IN APPENDIX B: http://www.doh.wa.gov/Tobacco/Data/County_Profiles/Profiles_2002/AppndxB.pdf

MAPS OF COUNTY RANKINGS: : http://www.doh.wa.gov/Tobacco/Data/County_Profiles/Profiles_2002/AppndxC.pdf

REFERENCES: http://www.doh.wa.gov/Tobacco/Data/County_Profiles/Profiles_2002/AppndxD.pdf